












エルエイジー10液の希釈について

吉田製薬株式会社
学術部

効能・効果		濃度 (塩酸アルキルジアミノ エチルグリシンとして)	希釈倍数	エルエイジー10液 使用量(全量1L)
生体	手指・皮膚の消毒 	0.05% ～ 0.2%	200倍 ～ 50倍	5mL ～ 20mL
	手術部位(手術野)の皮膚の消毒 	0.1% 0.2%	100倍 50倍	10mL 20mL
	手術部位(手術野)の粘膜の消毒 	0.01% ～ 0.05%	1000倍 ～ 200倍	1mL ～ 5mL
	皮膚の創傷部位の消毒 	0.01% ～ 0.05%	1000倍 ～ 200倍	1mL ～ 5mL
	粘膜の創傷部位の消毒 	0.01% ～ 0.05%	1000倍 ～ 200倍	1mL ～ 5mL
器具および環境	医療用具の消毒 	0.05% ～ 0.2%	200倍 ～ 50倍	5mL ～ 20mL
	結核領域における医療用具の消毒 	0.2% ～ 0.5%	50倍 ～ 20倍	20mL ～ 50mL
	手術室・病室・家具・器具・ 物品などの消毒  	0.05% ～ 0.2%	200倍 ～ 50倍	5mL ～ 20mL
	結核領域における手術室・病室 ・家具・器具・物品などの消毒  	0.2% ～ 0.5%	50倍 ～ 20倍	20mL ～ 50mL