











逆性石ケン液10「ヨシダ」の希釈について

吉田製薬株式会社
学術部

効能・効果		濃度 (塩化ベンザル コニウムとして)	希釈倍数	逆性石ケン液10 使用量(全量1L)
生 体	手指・皮膚の消毒 	0.05% ～ 0.1%	200倍 ～ 100倍	5mL ～ 10mL
	手術部位(手術野)の皮膚消毒 	0.1%5分洗後 0.2%塗布	100倍 50倍	10mL 20mL
	手術部位(手術野)の粘膜消毒 	0.01% ～ 0.025%	1000倍 ～ 400倍	1mL ～ 2.5mL
	皮膚の創傷部位の消毒 	0.01% ～ 0.025%	1000倍 ～ 400倍	1mL ～ 2.5mL
	粘膜の創傷部位の消毒 	0.01% ～ 0.025%	1000倍 ～ 400倍	1mL ～ 2.5mL
	感染皮膚面の消毒 	0.01%	1000倍	1mL
	腔の消毒 	0.02% ～ 0.05%	500倍 ～ 200倍	2mL ～ 5mL
	結膜囊の洗浄・消毒 	0.01% ～ 0.05%	1000倍 ～ 200倍	1mL ～ 5mL
器 具 お よ び 環 境	医療器具の消毒 	0.1%	100倍	10mL
	手術室・病室・家具・器具・ 物品などの消毒 	0.05% ～ 0.2%	200倍 ～ 50倍	5mL ～ 20mL